

Differential Contribution of L-, N-, and P/Q-type Calcium Channels to $[Ca^{2+}]_i$ Changes Evoked by Kainate in Hippocampal Neurons

Ana R. Santiago · Caetana M. Carvalho ·
Arsélio P. Carvalho · António F. Ambrósio

Accepted: 30 January 2008 / Published online: 27 March 2008
© Springer Science+Business Media, LLC 2008

Abstract We investigated the contribution of L-, N- and P/Q-type Ca^{2+} channels to the $[Ca^{2+}]_i$ changes, evoked by kainate, in the cell bodies of hippocampal neurons, using a pharmacological approach and Ca^{2+} imaging. Selective Ca^{2+} channel blockers, namely nitrendipine, ω -Conotoxin GVIA (ω -GVIA) and ω -Agatoxin IVA (ω -AgaIVA) were used. The $[Ca^{2+}]_i$ changes evoked by kainate presented a high variability, and were abolished by NBQX, a AMPA/kainate receptor antagonist, but the *N*-methyl-D-aspartate (NMDA) receptor antagonist, D-AP5, was without effect. Each Ca^{2+} channel blocker caused differential inhibitory effects on $[Ca^{2+}]_i$ responses evoked by kainate. We grouped the neurons for each blocker in three subpopulations: (1) neurons with responses below 60% of the control; (2) neurons with responses between 60% and 90% of the control, and (3) neurons with responses above 90% of the control. The inhibition caused by nitrendipine was higher than the inhibition caused by ω -GVIA or ω -AgaIVA. Thus, in the presence of nitrendipine, the percentage of cells with responses below 60% of the control was 41%, whereas in the case of ω -GVIA or ω -AgaIVA the values were 9 or 17%, respectively. The results indicate that hippocampal neurons differ in what concerns their L-, N- and P/Q- type

Ca^{2+} channels activated by stimulation of the AMPA/kainate receptors.

Keywords Voltage-sensitive calcium channels · $[Ca^{2+}]_i$ changes · AMPA and kainate receptors · Hippocampal neurons

Introduction

Ionotropic glutamate receptors are ligand-gated ion channels, permeable to cations, and have been classified into three subtypes: *N*-methyl-D-aspartate (NMDA), α -amino-3-hydroxy-5-methyl-4-isoxazole propionic acid (AMPA) and kainate receptors. NMDA receptors are responsible for the slow component of excitatory postsynaptic potentials, whereas AMPA and kainate receptors mediate fast excitatory transmission. These receptors are involved in important physiological and pathological processes, such as cell differentiation and growing, learning and memory, and excitotoxicity [1].

The activation of ionotropic glutamate receptors causes a substantial Na^+ entry into the cell, and the subsequent cell depolarisation triggers the opening of voltage-sensitive Ca^{2+} channels (VSCCs) and Ca^{2+} entry, which may also occur directly through NMDA receptors, as well as through Ca^{2+} permeable AMPA and kainate receptors, depending on their subunit composition.

In nerve cells, L-type Ca^{2+} channels are predominantly located in cell bodies and dendrites [2–4]. N-type Ca^{2+} channels are mainly located in nerve terminals and dendrites, although they can also be found in cell bodies [5–7]. P-type and Q-type Ca^{2+} channels are predominantly located in nerve terminals, but they may also be present in dendrites and cell bodies [2, 8–10]. The pharmacological

Special issue article in honor of Dr. Ricardo Tapia.

A. R. Santiago · C. M. Carvalho · A. P. Carvalho (✉) ·
A. F. Ambrósio

Center for Neuroscience and Cell Biology, Department of
Zoology, University of Coimbra, Coimbra 3004-517, Portugal
e-mail: carvalho@cnc.uc.pt

A. F. Ambrósio
Center of Ophthalmology, Institute of Biomedical Research on
Light and Image (IBILI), Faculty of Medicine, University of
Coimbra, Coimbra, Portugal

distinction between P- and Q-type Ca^{2+} channels is difficult because they are both composed by the α_{1A} subunit, and therefore they are usually named P/Q-type Ca^{2+} channels [11, 12].

We have previously shown that VSCCs, mainly L- and N-type Ca^{2+} channels, have a predominant role in the $[\text{Ca}^{2+}]_i$ changes evoked by kainate in cultured hippocampal neurons measured in cell populations [13]. The $[\text{Ca}^{2+}]_i$ responses evoked by kainate have a high degree of variability among individual hippocampal neurons [14], which is likely due to a differential expression of AMPA and kainate receptors. These variations might have consequences in synaptic efficacy and information processing. However, the variability in $[\text{Ca}^{2+}]_i$ responses may be also due to a differential distribution of VSCCs among hippocampal neurons. In the present work we investigated and quantified the contribution of several Ca^{2+} channels to the Ca^{2+} transients induced by AMPA/kainate receptor activation in single hippocampal neurons, by Ca^{2+} imaging.

Experimental Procedure

Materials

Neurobasal medium, B27 supplement, gentamicin and trypsin (USP grade) were purchased from GIBCO BRL, Life Technologies, Scotland. DNase (DN-25) was purchased from Sigma Chemical, St. Louis, MO, USA. Kainate, 1,2,3,4-Tetrahydro-6-nitro-2,3-dioxo-benzof[quinoxaline-7-sulfonamide (NBQX) and D(–)-2-amino-5-phosphonopentanoic acid (D-AP5) were purchased from TOCRIS, Bristol, UK. The acetoxymethyl ester of Fura-2 (Fura-2/AM) and Pluronic F-127 were purchased from Molecular Probes, Leiden, The Netherlands. Nitrendipine was a kind gift of Dr G. Terstappen, from Bayer AG, Germany. ω -GVIA was obtained from Peninsula Laboratories, Belmont, CA, USA, and ω -AgaIVA was from Peptide Institute, Osaka, Japan. All other reagents were from Sigma Chemical, St. Louis, MO, USA or from Merck-Schuchardt, Germany.

Cell Culture

All procedures involving animals were conducted in accordance with the “Principles of laboratory animal care” (NIH publication No. 86-23, revised 1985) statement for the use of animals.

Hippocampal neurons were obtained from the hippocampi of E18-E19 Wistar rat embryos, after treatment with trypsin (1.0 mg/ml; 15 min; 37°C) and deoxyribonuclease I (0.15 mg/ml) in Ca^{2+} - and Mg^{2+} -free Hank's balanced salt solution (137 mM NaCl, 5.36 mM KCl, 0.44 mM KH_2PO_4 , 0.34 mM $\text{Na}_2\text{HPO}_4 \cdot 2\text{H}_2\text{O}$, 4.16 mM NaHCO_3 ,

5 mM glucose, supplemented with 0.001% phenol red, 1 mM sodium pyruvate, 10 mM HEPES, pH 7.4). Hippocampal neurons were cultured in B27-supplemented Neurobasal medium (GIBCO), a serum-free medium combination [15], supplemented with glutamate (25 μM), glutamine (0.5 mM) and gentamicin (0.12 mg/ml). Cultures were kept at 37°C in a humidified incubator in 5% CO_2 /95% air, for 7–8 days, the time required for maturation of hippocampal neurons.

The cells were plated at a density of 45×10^3 cells/cm² on poly-D-lysine-coated coverslips.

Fura-2 Fluorescence Imaging in Individual Cells

The changes in intracellular free Ca^{2+} concentration $[\text{Ca}^{2+}]_i$ of individual cells were measured using the fluorescent probe Fura-2 [16]. Hippocampal neurons were incubated with 5 μM Fura-2/AM (Molecular Probes) and 0.02% Pluronic F-127 (Molecular Probes) for 30 min at 37°C, in Krebs buffer (142 mM NaCl, 4 mM KCl, 1 mM MgCl_2 , 1 mM CaCl_2 , 10 mM glucose, 10 mM Hepes-Na, pH 7.4) supplemented with 0.1% bovine serum albumine (BSA; Calbiochem-Boehringer). After incubation, the coverslips were rinsed with Krebs buffer and placed in a perfusion chamber on the stage of an inverted Nikon Diaphot fluorescence microscope.

The experiments were conducted under continuous perfusion with warm Krebs buffer (30°C) with or without drugs, as indicated in the figure legends. Each recording lasted 75 s, and neurons were stimulated 20 s after image acquisition. The fluorescence changes were recorded with a multiple excitation Magical imaging system (Applied Imaging, UK). Hippocampal neurons were alternately excited at 340 and 380 nm using a switching filter wheel, and the emitted fluorescence, collected with a 40 \times objective (Nikon) was driven to a SIT camera (Silicon Intensified Target; Photonics Science), after passing through a 510 nm bandpass filter. Image analysis was performed with the Magical imaging system. Briefly, the background fluorescence at each wavelength (340 and 380 nm) was subtracted and the ratio of the fluorescence images on a pixel-by-pixel basis was obtained. The data were stored as 8-bit pseudocolor images. Areas of the cell bodies were drawn and the averaged value of pixel intensities was evaluated at each time point, in order to obtain fluorescence ratio versus time plots for all areas defined. The results were expressed as the ratio of fluorescence intensity with excitation at 340 and 380 nm versus time.

Design of Ca^{2+} Experiments

Each experiment consists of two independent recordings, and the first recording was considered an internal control for each cell. The hippocampal neurons were stimulated

twice with kainate ($t = 20$ s), which was present until the end of each recording ($t = 75$ s). The cells were allowed to recover for 15 min between the first and second stimulus, and each drug tested was present 5 min before the second stimulus, until the end of the experiment.

Data Analysis

The data are expressed as mean \pm SEM. Significance was considered at $P < 0.05$ using a two sample Student's t test.

Results

Characterization of the $[Ca^{2+}]_i$ Responses Evoked by Kainate in Hippocampal Neurons

We analyzed the $[Ca^{2+}]_i$ changes evoked by 100 μ M kainate in the cell bodies. As expected, there was a high variability among cells in terms of $[Ca^{2+}]_i$ changes. We considered the $[Ca^{2+}]_i$ changes as the differences between the peak, after stimulus, and the basal value of the 340 nm/380 nm fluorescence ratio. In all experiments, the first stimulus was used as an internal control.

In control experiments, hippocampal neurons were stimulated twice with kainate, in the absence of calcium channel blockers. The second $[Ca^{2+}]_i$ response was similar to the first one ($95.8 \pm 1.5\%$ of the control; Fig. 1a) in all hippocampal neurons analyzed ($n = 48$). The AMPA/kainate receptor antagonist, NBQX (10 μ M), abolished the $[Ca^{2+}]_i$ changes (Fig. 1b). The presence of D-AP5 (100 μ M), a NMDA receptor antagonist, did not cause any inhibitory effect on the $[Ca^{2+}]_i$ changes ($97.0 \pm 1.3\%$ of the control; $n = 31$ neurons), thus excluding NMDA receptor involvement in the $[Ca^{2+}]_i$ response to kainate.

Contribution of L-type Ca^{2+} Channels to the $[Ca^{2+}]_i$ Changes Evoked by Kainate

The contribution of L-type Ca^{2+} channels to the $[Ca^{2+}]_i$ changes evoked by kainate was analyzed by stimulating neurons in the presence of nitrendipine (10 μ M). The L-type Ca^{2+} channel blocker, nitrendipine, inhibited the $[Ca^{2+}]_i$ changes to $60.5 \pm 1.9\%$ of the control ($n = 58$ cells; $P < 0.05$). Since the inhibitory effect of nitrendipine




Fig. 1 Representative recordings of kainate-evoked $[Ca^{2+}]_i$ changes, measured with the fluorescent probe Fura-2, in single hippocampal neurons. **(A)** Control; **(B)** Effect of NBQX (antagonist of AMPA/kainate receptors). The results are expressed as the ratio of fluorescence intensity with excitation at 340 and 380 nm (emission at 510 nm) versus time. Each experiment consists of two independent recordings (75 s each), and the first recording was considered an internal control for each cell. The $[Ca^{2+}]_i$ changes were evoked by 100 μ M kainate, and were analyzed in cell bodies. The cells were stimulated twice with kainate ($t = 20$ s), with a 15 min recovery interval between the first and the second stimulus. NBQX was present 5 min before the second stimulus, until the end of the experiment. **(A)** In control experiments, hippocampal neurons were stimulated twice with kainate, with no drug present during the second stimulus. The second recording was similar to the first one in all 48 cells analyzed from three independent cell cultures. **(B)** NBQX abolished the kainate evoked- $[Ca^{2+}]_i$ changes in the 38 cells analyzed from two independent cell cultures

was very heterogeneous in the cell population, the cells were grouped in three subpopulations, as follows: responses to kainate in the presence of nitrendipine below 60% of the control (subpopulation 1), between 60% and 90% of the control (subpopulation 2), and above 90% of the control (subpopulation 3) (Table 1). Considering the first group

Table 1 Effect of nitrendipine on $[Ca^{2+}]_i$ changes evoked by kainate in the total population, and in subpopulations 1, 2 and 3

$\Delta [Ca^{2+}]_i$ (% of control \pm SEM)			
Total population	Subpopulation 1 (<60%)	Subpopulation 2 (60–90%)	Subpopulation 3 (>90%)
$60.5 \pm 1.9\%$ (100%)	$46.2 \pm 1.9\%$ (41.4%)	$69.9 \pm 1.1\%$ (56.9%)	94% (1.7%)

The percentage of hippocampal neurons belonging to each population is indicated in parenthesis

of cells, responses below 60% of the control (first stimulus), the $[Ca^{2+}]_i$ response evoked by kainate, in the presence of nitrendipine, was $46.2 \pm 1.9\%$ of the control (Fig. 2a).




Fig. 2 Effect of nitrendipine (L-type Ca^{2+} channel blocker) on the $[Ca^{2+}]_i$ changes evoked by kainate ($100 \mu M$) in single hippocampal neurons. The results are expressed as the ratio of fluorescence intensity with excitation at 340 and 380 nm (emission at 510 nm) versus time. Each experiment consists of two independent recordings (75 s each), and the first recording was considered an internal control for each cell. The $[Ca^{2+}]_i$ changes were considered as the differences between the peak, after stimulus, and the basal value of the 340 nm/380 nm fluorescence ratio. The $[Ca^{2+}]_i$ changes were analyzed in cell bodies. Nitrendipine (Nit; $10 \mu M$) was present 5 min before the second stimulus, until the end of each experiment. Since the inhibitory effect of nitrendipine was very heterogeneous, the cells were grouped in three subpopulations. The recordings shown above are representative of each cell population, with $[Ca^{2+}]_i$ responses below 60% of the control (A), between 60% and 90% of the control (B), and above 90% of the control (C). A total of 58 cells, from four independent cell cultures, were analyzed, and the quantification of the data is shown in Table 1

This group of cells represents 41.4% of the cells stimulated in the presence of nitrendipine. In the second group of cells (responses between 60% and 90% of the control), which represents 56.9% of cells (33 neurons), nitrendipine inhibited the $[Ca^{2+}]_i$ increase to $69.9 \pm 1.1\%$ of the control (Fig. 2b). The $[Ca^{2+}]_i$ response was above 90% of the control (94%) in just one cell (Fig. 2c), thus indicating that L-type Ca^{2+} channels have a major contribution to the $[Ca^{2+}]_i$ changes upon non-NMDA glutamate receptors activation in the majority of hippocampal neurons.

Contribution of N- and P/Q-type Ca^{2+} Channels to the $[Ca^{2+}]_i$ Changes Evoked by Kainate

The contribution of N- and P/Q-type Ca^{2+} channels to the $[Ca^{2+}]_i$ changes evoked by kainate were also investigated, by stimulating the cells in the presence of ω -GVIA or ω -AgaIVA, respectively. The $[Ca^{2+}]_i$ changes were inhibited $20.1 \pm 1.7\%$ ($P < 0.05$) or $17.2 \pm 2.2\%$ ($P < 0.05$) when neurons were stimulated in the presence of ω -GVIA (500 nM) or ω -AgaIVA (100 nM), respectively. Again, due to the variability of the inhibitory effects caused by these Ca^{2+} channel blockers, the cells were grouped in three different populations, as indicated in Tables 2 and 3.

Relatively to the contribution of N-type Ca^{2+} channels to the $[Ca^{2+}]_i$ changes, 76 neurons were analyzed. In a small percentage of neurons (9.2%; seven cells), the $[Ca^{2+}]_i$ changes evoked by kainate, with ω -GVIA present, were below 60% of the control ($48.1 \pm 3.7\%$ of the control; Fig. 3a). In the second population of neurons ($[Ca^{2+}]_i$ responses between 60% and 90% of the control), which represents the majority of neurons (73.7% of neurons), ω -GVIA inhibited the $[Ca^{2+}]_i$ increase to $79.6 \pm 0.9\%$ of the control (Fig. 3b; Table 2). In 13 neurons (17.1% of neurons) the responses were not significantly affected by the presence of ω -GVIA ($98.4 \pm 1.5\%$ of the control; Fig. 3c).

The results obtained when neurons were stimulated in the presence of the P/Q-type Ca^{2+} channel blocker, ω -AgaIVA, were similar to those obtained with ω -GVIA. In these experiments, 48 neurons were analyzed. Considering the first population (neurons with responses below 60% of the control), the presence of ω -AgaIVA inhibited the $[Ca^{2+}]_i$ increase to $51.7 \pm 2.9\%$ of the control (Fig. 4a), corresponding to 16.7% of neurons (eight cells). In the second population, that includes the majority of neurons (66.7% of cells), the $[Ca^{2+}]_i$ increase was inhibited to $72.3 \pm 1.5\%$ of the control (Fig. 4b). In the third population (eight neurons; 16.6% of cells), ω -AgaIVA was without effect, and the $[Ca^{2+}]_i$ response was $95.8 \pm 1.8\%$ of the control (Fig. 4c; Table 3).

Table 2 Effect of ω -GVIA on $[Ca^{2+}]_i$ changes evoked by kainate in the total population, and in subpopulations 1, 2 and 3

$\Delta [Ca^{2+}]_i$ (% of control \pm S.E.M.)			
Total population	Subpopulation 1 (<60%)	Subpopulation 2 (60–90%)	Subpopulation 3 (>90%)
79.9 \pm 1.7% (100%)	48.1 \pm 3.7% (9.2 %)	79.6 \pm 0.9% (73.7 %)	98.4 \pm 1.5% (17.1%)

The percentage of hippocampal neurons belonging to each population is indicated in parenthesis

Table 3 Effect of ω -AgaIVA on $[Ca^{2+}]_i$ changes evoked by kainate in the total population, and in subpopulations 1, 2 and 3

$\Delta [Ca^{2+}]_i$ (% of control \pm SEM)			
Total population	Subpopulation 1 (<60%)	Subpopulation 2 (60–90%)	Subpopulation 3 (>90%)
82.8 \pm 2.2% (100%)	51.7 \pm 2.9% (16.7%)	72.3 \pm 1.5% (66.7%)	95.8 \pm 1.8% (16.6%)

The percentage of hippocampal neurons belonging to each population is indicated in parenthesis

Discussion

We have previously shown that VSCCs, particularly L- and N-type Ca^{2+} channels, have a large contribution to the $[Ca^{2+}]_i$ changes evoked by kainate, measured in populations of cultured hippocampal neurons [13]. We also demonstrated that the alterations in the $[Ca^{2+}]_i$ evoked by kainate or AMPA, in individual neurons, present a high cell-to-cell variability, and, in addition, hippocampal neurons have differential sensitivities to LY 303070 and cyclothiazide, which are, respectively, an antagonist and an allosteric modulator of AMPA receptors [14]. Those observations suggest that there is a differential expression and localization of AMPA receptors in hippocampal neurons. Since VSCCs are important mediators of glutamatergic neurotransmission and contribute to $[Ca^{2+}]_i$ changes upon glutamate receptor activation, in the present work we investigated whether different types of VSCCs are differentially coupled to the activation of AMPA and kainate receptors, in single hippocampal neurons.

As previously demonstrated, there was a high cell-to-cell variability in the $[Ca^{2+}]_i$ changes evoked by kainate, again indicating that AMPA and kainate receptors are differentially distributed in hippocampal neurons. The $[Ca^{2+}]_i$ changes were due to AMPA/kainate receptor activation, since NBQX, the AMPA receptor antagonist, abolished the $[Ca^{2+}]_i$ responses, which were not affected by the presence of D-AP5, the NMDA receptor antagonist. Endogenous glutamate, released upon AMPA/kainate receptor activation, could activate NMDA receptors, which might in turn further contribute to the Ca^{2+} signal. However, the blockade of NMDA receptors did not inhibit the $[Ca^{2+}]_i$ changes, indicating that NMDA receptors do not contribute to the $[Ca^{2+}]_i$ changes in these conditions, as was shown previously with $^{45}Ca^{2+}$ uptake experiments and AMPA receptor stimulation in hippocampal neurons [17].

L-type Ca^{2+} channels are particularly abundant in cell bodies and dendrites [3, 18], and several reports have shown that L-type Ca^{2+} channels are activated upon cell stimulation with kainate [19–21]. The present results confirm that these Ca^{2+} channels have an important role in coupling AMPA/kainate receptor activation with the Ca^{2+} signal. Indeed, considering the entire population studied, the inhibitory effect caused by nitrendipine (39.5%) was much higher than that caused either by ω -CgTx GVIA (20.1%) or ω -Aga IVA (17.2%). In addition, the $[Ca^{2+}]_i$ responses evoked by kainate, in the presence of nitrendipine, decreased below 60% of the control in about 40% of neurons. In the presence of either ω -CgTx GVIA or ω -Aga IVA, the $[Ca^{2+}]_i$ responses decreased below 60% of the control in only 9 or 17% of neurons, respectively, much lower than 40% observed for nitrendipine. These facts clearly demonstrate that L-type Ca^{2+} channels have a predominant role in the $[Ca^{2+}]_i$ transients evoked by kainate, in the cell bodies of hippocampal neurons.

It has been demonstrated that N- and P/Q-type Ca^{2+} channels are predominantly located in dendrites and nerve terminals, where they are closely associated with the release machinery [9, 22, 23]. The blockade of these channels has been shown to inhibit the release of several neurotransmitters [24–27] and inhibits glutamatergic synaptic transmission in the hippocampus [23, 28]. We have also previously shown that these channels have a major contribution to the $[Ca^{2+}]_i$ signal in isolated hippocampal nerve terminals [29]. However, in these cultures, N- and P/Q-type Ca^{2+} channels also contributed to the $[Ca^{2+}]_i$ increase in cell bodies, in several neurons, since the $[Ca^{2+}]_i$ responses were inhibited either by ω -CgTx GVIA or ω -Aga IVA in those neurons. These observations clearly indicate that these channels are located in cell bodies and dendrites, at least in some hippocampal neurons. Indeed, it has been demonstrated that both N- and P/Q-type Ca^{2+}




Fig. 3 Effect of ω -GVIA (N-type Ca^{2+} channel blocker) on the $[\text{Ca}^{2+}]_i$ changes evoked by kainate (100 μM) in single hippocampal neurons. The results are expressed as the ratio of fluorescence intensity with excitation at 340 and 380 nm (emission at 510 nm) versus time. Each experiment consists of two independent recordings (75 s each), and the first recording was considered an internal control for each cell. The $[\text{Ca}^{2+}]_i$ changes were considered as the differences between the peak, after stimulus, and the basal value of the 340 nm/380 nm fluorescence ratio. The $[\text{Ca}^{2+}]_i$ changes were analyzed in cell bodies. ω -GVIA (500 nM) was present 5 min before the second stimulus, until the end of each experiment. Since the inhibitory effect of ω -GVIA was heterogeneous, the cells were separated in three subpopulations. The recordings above are representative of each cell population with $[\text{Ca}^{2+}]_i$ responses below 60% of the control (**A**), between 60% and 90% of the control (**B**), and above 90% of the control (**C**). A total of 76 neurons, from three independent cell cultures, were analyzed, and the quantification of the data is shown in Table 2




Fig. 4 Effect of ω -AgaIVA (P/Q-type Ca^{2+} channel blocker) on the $[\text{Ca}^{2+}]_i$ changes evoked by kainate (100 μM) in single hippocampal neurons. The results are expressed as the ratio of fluorescence intensity with excitation at 340 and 380 nm (emission at 510 nm) versus time. Each experiment consists of two independent recordings (75 s each), and the first recording was considered an internal control for each cell. The $[\text{Ca}^{2+}]_i$ changes were considered as the differences between the peak, after stimulus, and the basal value of the 340 nm/380 nm fluorescence ratio. The $[\text{Ca}^{2+}]_i$ changes were analyzed in cell bodies. ω -AgaIVA (100 nM) was present 5 min before the second stimulus, until the end of each experiment. Since the inhibitory effect of ω -AgaIVA was heterogeneous, the cells were separated in three subpopulations. The recordings above are representative of each cell population, with $[\text{Ca}^{2+}]_i$ responses below 60% of the control (**A**), between 60% and 90% of the control (**B**), and above 90% of the control (**C**). A total of 48 neurons, from three independent cell cultures, were analyzed, and the quantification of the data is shown in Table 3

channels are located in cell bodies and dendrites of several types of neurons, including hippocampal neurons, in addition to their location in nerve terminals [6, 8, 18,

30–32]. Also, these channels contribute to the stimulus-induced postsynaptic Ca^{2+} fluxes in hippocampal neurons [33], and N-type Ca^{2+} channels are primarily involved in

the $[Ca^{2+}]_i$ increase evoked by high potassium in the pyramidal neurons of the dorsal cochlear nucleus [34]. This group of findings indicates that, at least in some cells, N- and P/Q-type Ca^{2+} channels have an important role in glutamatergic synaptic transmission, not only because they regulate transmitter release from nerve terminals, as previously demonstrated, but also because they mediate Ca^{2+} entry in postsynaptic neurons upon the activation of ionotropic glutamate receptors.

In the present work, we showed that the contribution of different types of VSCCs to the $[Ca^{2+}]_i$ signal evoked by kainate is very heterogeneous. In a few cells, the $[Ca^{2+}]_i$ changes were inhibited by about 60% either by nitrendipine, ω -CgTx GVIA or ω -Aga IVA, and in others these blockers were without effect. In some of those cells, either L-, N-, or P/Q-type Ca^{2+} channels are clearly important in coupling glutamate receptor activation with Ca^{2+} signal, probably because these Ca^{2+} channels are particularly abundant in cell bodies and dendrites of those cells or also because both AMPA/kainate receptors and one certain type of Ca^{2+} channels are located in near proximity. In other cells, N- and P/Q-type Ca^{2+} channels did not contribute to the $[Ca^{2+}]_i$ changes or had a low contribution, likely because these Ca^{2+} channels are not located or their expression is rather low in the cell bodies and dendrites of those neurons.

In summary, our results show that the activation of L-type Ca^{2+} channels is coupled to non-NMDA glutamate receptor activation in the cell bodies of hippocampal neurons, and these channels are also a major route for Ca^{2+} entry, although its contribution varies from cell to cell. Regarding N- or P/Q-type Ca^{2+} channels, the results indicate that they are not present in the cell bodies or dendrites of all hippocampal neurons. In some cells, they have an important role in coupling AMPA/kainate receptor activation to Ca^{2+} increase, but in others these Ca^{2+} channels do not contribute at all to Ca^{2+} entry into the cell bodies. The fact that different types of Ca^{2+} channels are differentially coupled to the activation of non-NMDA receptors will certainly have consequences in terms of information processing in neurons and will contribute to variations in cell responses in hippocampal and brain circuitries.

Acknowledgement This work was supported by Fundação para a Ciência e a Tecnologia (FCT), Portugal (Grant POCTI/NSE/35875/2000).

References

- Barnes GN, Slevin JT (2003) Ionotropic glutamate receptor biology: effect on synaptic connectivity and function in neurological disease. *Curr Med Chem* 10:2059–2072
- Elliott EM, Malouf AT, Catterall WA (1995) Role of calcium channel subtypes in calcium transients in hippocampal CA3 neurons. *J Neurosci* 15:6433–6444
- Timmermann DB, Westenbroek RE, Schousboe A, Catterall WA (2002) Distribution of high-voltage-activated calcium channels in cultured gamma-aminobutyric acidergic neurons from mouse cerebral cortex. *J Neurosci Res* 67:48–61
- Westenbroek RE, Ahljianian MK, Catterall WA (1990) Clustering of L-type Ca^{2+} channels at the base of major dendrites in hippocampal pyramidal neurons. *Nature* 347:281–284
- Reynolds JJ, Wagner JA, Snyder SH, Thayer SA, Olivera BM, Miller RJ (1986) Brain voltage-sensitive calcium channel subtypes differentiated by omega-conotoxin fraction GVIA. *Proc Natl Acad Sci USA* 83:8804–8807
- Westenbroek RE, Hell JW, Warner C, Dubel SJ, Snutch TP, Catterall WA (1992) Biochemical properties and subcellular distribution of an N-type calcium channel alpha 1 subunit. *Neuron* 9:1099–1115
- Westenbroek RE, Sakurai T, Elliott EM et al (1995) Immunohistochemical identification and subcellular distribution of the alpha 1A subunits of brain calcium channels. *J Neurosci* 15:6403–6418
- Nakanishi S, Fujii A, Kimura T, Sakakibara S, Mikoshiba K (1995) Spatial distribution of omega-agatoxin IVA binding sites in mouse brain slices. *J Neurosci Res* 41:532–539
- Qian J, Noebels JL (2001) Presynaptic Ca^{2+} channels and neurotransmitter release at the terminal of a mouse cortical neuron. *J Neurosci* 21:3721–3728
- Takahashi T, Momiyama A (1993) Different types of calcium channels mediate central synaptic transmission. *Nature* 366:156–158
- Sather WA, Tanabe T, Zhang JF, Mori Y, Adams ME, Tsien RW (1993) Distinctive biophysical and pharmacological properties of class A (BI) calcium channel alpha 1 subunits. *Neuron* 11:291–303
- Stea A, Tomlinson WJ, Soong TW et al (1994) Localization and functional properties of a rat brain alpha 1A calcium channel reflect similarities to neuronal Q- and P-type channels. *Proc Natl Acad Sci USA* 91:10576–10580
- Ambrosio AF, Silva AP, Malva JO, Soares-da-Silva P, Carvalho AP, Carvalho CM (1999) Carbamazepine inhibits L-type Ca^{2+} channels in cultured rat hippocampal neurons stimulated with glutamate receptor agonists. *Neuropharmacology* 38:1349–1359
- Ambrosio AF, Silva AP, Malva JO, Mesquita JF, Carvalho AP, Carvalho CM (2000) Role of desensitization of AMPA receptors on the neuronal viability and on the $[Ca^{2+}]_i$ changes in cultured rat hippocampal neurons. *Eur J Neurosci* 12:2021–2031
- Brewer GJ, Torricelli JR, Evege EK, Price PJ (1993) Optimized survival of hippocampal neurons in B27-supplemented Neurobasal, a new serum-free medium combination. *J Neurosci Res* 35:567–576
- Gryniewicz G, Poenie M, Tsien RY (1985) A new generation of Ca^{2+} indicators with greatly improved fluorescence properties. *J Biol Chem* 260:3440–3450
- Araujo IM, Carreira BP, Pereira T et al (2007) Changes in calcium dynamics following the reversal of the sodium-calcium exchanger have a key role in AMPA receptor-mediated neurodegeneration via calpain activation in hippocampal neurons. *Cell Death Differ* 14:1635–1646
- Pravettoni E, Bacci A, Coco S, Forbicini P, Matteoli M, Verderio C (2000) Different localizations and functions of L-type and N-type calcium channels during development of hippocampal neurons. *Dev Biol* 227:581–594
- Alberdi E, Sanchez-Gomez MV, Marino A, Matute C (2002) Ca^{2+} influx through AMPA or kainate receptors alone is sufficient to initiate excitotoxicity in cultured oligodendrocytes. *Neurobiol Dis* 9:234–243
- Courtney MJ, Lambert JJ, Nicholls DG (1990) The interactions between plasma membrane depolarization and glutamate receptor activation in the regulation of cytoplasmic free calcium in cultured cerebellar granule cells. *J Neurosci* 10:3873–3879

21. Leski ML, Valentine SL, Coyle JT (1999) L-type voltage-gated calcium channels modulate kainic acid neurotoxicity in cerebellar granule cells. *Brain Res* 828:27–40
22. Scholz KP, Miller RJ (1995) Developmental changes in presynaptic calcium channels coupled to glutamate release in cultured rat hippocampal neurons. *J Neurosci* 15:4612–4617
23. Wheeler DB, Randall A, Tsien RW (1994) Roles of N-type and Q-type Ca^{2+} channels in supporting hippocampal synaptic transmission. *Science* 264:107–111
24. Burke SP, Adams ME, Taylor CP (1993) Inhibition of endogenous glutamate release from hippocampal tissue by Ca^{2+} channel toxins. *Eur J Pharmacol* 238:383–386
25. Keith RA, Mangano TJ, Salama AI (1989) Inhibition of N-methyl-D-aspartate- and kainic acid-induced neurotransmitter release by omega-conotoxin GVIA. *Br J Pharmacol* 98:767–772
26. Kimura M, Yamanishi Y, Hanada T et al (1995) Involvement of P-type calcium channels in high potassium-elicited release of neurotransmitters from rat brain slices. *Neuroscience* 66:609–615
27. Saydoff JA, Zaczek R (1996) Blockade of N- and Q-type Ca^{2+} channels inhibit K^{+} -evoked $[\text{H}^3]\text{acetylcholine}$ release in rat hippocampal slices. *Brain Res Bull* 40:283–286
28. Luebke JI, Dunlap K, Turner TJ (1993) Multiple calcium channel types control glutamatergic synaptic transmission in the hippocampus. *Neuron* 11:895–902
29. Ambrosio AF, Malva JO, Carvalho AP, Carvalho CM (1997) Inhibition of N-,P/Q- and other types of Ca^{2+} channels in rat hippocampal nerve terminals by the adenosine A1 receptor. *Eur J Pharmacol* 340:301–310
30. Mills LR, Niesen CE, So AP, Carlen PL, Spigelman I, Jones OT (1994) N-type Ca^{2+} channels are located on somata, dendrites, and a subpopulation of dendritic spines on live hippocampal pyramidal neurons. *J Neurosci* 14:6815–6824
31. Wang G, Ding S, Yunokuchi K (2000) Different distribution of nifedipine- and omega-conotoxin GVIA-sensitive Ca^{2+} channels in rat hippocampal neurons. *Neuroreport* 11:2419–2423
32. Wang G, Hashiguchi T, Sakamoto Y, Takigawa M, Kameyama M (1999) Distribution of nifedipine- and omega-conotoxin GVIA-sensitive Ca^{2+} channels in cultured rat neocortical neurons. *Neuroscience* 93:491–496
33. Igelmund P, Zhao YQ, Heinemann U (1996) Effects of T-type, L-type, N-type, P-type, and Q-type calcium channel blockers on stimulus-induced pre- and postsynaptic calcium fluxes in rat hippocampal slices. *Exp Brain Res* 109:22–32
34. Rusznak Z, Harasztosi C, Stanfield PR, Kovacs L, Szucs G (2000) Potassium-depolarization-induced cytoplasmic $[\text{Ca}^{2+}]$ transient in freshly dissociated pyramidal neurones of the rat dorsal cochlear nucleus. *Pflugers Arch* 440:462–466